

**Recombinant Human Apolipoprotein-Serum  
Amyloid A1  
(rHuApo-SAA1)**  
**PrimeGene Technical Data Sheet**

---

<b>Catalog Number:</b>	602-26
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 11.7 kDa, a single non-glycosylated polypeptide chain containing 104 amino acids.
<b>Quantity:</b>	10µg/50µg/1000µg
<b>AA Sequence:</b>	RSFFSFLGEA FDGARDMWRA YSDMREANYI GSDKYFHARG NYDAAKRGPG GVWAAEAISD ARENIQRFFG HGAEDSLADQ AANEWGRSGK DPNHFRPAGL PEKY
<b>Purity:</b>	> 98 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The biological activity determined by a chemoattract bioassay using human monocytes is in a concentration range of 10-100 ng/ml.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM Tris-HCl, pH 9.0, 150 mM NaCl.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

***Human Apolipoprotein-Serum Amyloid A1***

Serum amyloid A-1 protein which belongs to the SAA family is encoded by the SAA1 gene in humans. It is expressed by the liver and secreted in plasma. Serum amyloid A-1 is the major acute phase reactant and apolipoprotein of the HDL complex. Elevated serum SAA1 protein levels may be associated with lung cancer. The level of Apo-SAA1, normally 1-5 µg/ml in plasma, increases 500-1000 fold within 24 hours of an inflammatory stimulus and, under these conditions, is the most abundant HDL Apo-lipoprotein. Recombinant human Apo-SAA1 is an 11.7 kDa protein containing 104 amino acid residues.