

**Recombinant Human Insulin-like Growth
Factor-1 R36Q
(rHuIGF-1 R36Q)
PrimeGene Technical Data Sheet**

Catalog Number:	105-01V2
Source:	<i>E. coli</i>
Molecular Weight:	Approximately 7.7 kDa, a single polypeptide chain containing 70 amino acids.
Quantity:	25µg/100µg/1mg
AA Sequence:	GPETLCGAEL VDALQFVCGD RGFYFNKPTG YGSSSQRAPQ TGIVDECCFR SCDLRRLEMY CAPLKPAKSA
Purity:	> 95% by SDS-PAGE analyses.
Biological Activity:	Test in process.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, 150 mM NaCl, pH5.4.
Endotoxin:	Less than 0.1 EU/µg of rHuIGF-1 R36Q as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Insulin-like Growth Factor-1 R36Q

IGF-1 belonged to the insulin gene family, is a mitogenic polypeptide growth factor that stimulate the proliferation and survival of various cell types including muscle, bone, and cartilage tissue in vitro. It is produced primarily by the liver as an endocrine hormone as well as in target tissues in a paracrine/autocrine fashion. The production of IGF-1 is stimulated by growth hormone (GH) and can be retarded by undernutrition, growth hormone insensitivity, lack of growth hormone receptors, or failures of the downstream signaling pathway post GH receptor including SHP2 and STAT5B.