

**PrimeGene Technical Data Sheet**

---

<b>Catalog Number:</b>	101-17F
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 30.1 kDa, a disulfide-linked homodimer of two 134 amino acid polypeptide chains.
<b>Quantity:</b>	5µg/25µg/1000µg
<b>AA Sequence:</b>	MRKIPKVGHT FFQKPESCPP VPGGSMKLDI GIINENQRVS MSRNIESRST SPWNYTVTWD PNRYLPSEVVQ AQCRLGGIN AQQKEDISMN SVPIQQETLV VRRKHQGCSV SFQLEKVLVT VGCTCVTPVI HHVQ
<b>Purity:</b>	> 95 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by inducing IL-6 secretion of murine NIH/3T3 cells is less than 20 ng/ml, corresponding to a specific activity of > 5.0 × 10 <sup>4</sup> IU/mg.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
<b>Endotoxin:</b>	Less than 1 EU/µg of rHuIL-17F as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 6 mM HCl to a concentration of 0.1mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

***Human Interleukin-17F***

Human Interleukin-17F (IL-17F) is encoded by the IL17F gene located on the chromosome 6 and belongs to the IL-17 family which contains IL-17A, IL-17B, IL-17C, IL-17D, IL-17E and IL-17F. IL-17F that shares 50 % homologous of crystal structure to IL-17A and is expressed by activated T cells, has the functions of stimulating production of other cytokines (such as IL-6, IL-8 and granulocyte colony-stimulating factor) and proliferation of PBMC and T-cell, regulating cartilage matrix turnover, and inhibiting angiogenesis.