

Recombinant Murine Persephin (rMuPersephin)

PrimeGene Technical Data Sheet

Catalog Number:	127-23
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 20.7 kDa, a disulfide-linked homodimer of two 96 amino acid polypeptide chains.
Quantity:	5µg/20µg/1000µg
AA Sequence:	ALAGSCLWS LTLVPAELGL GYASEEKVIF RYCAGSCPQE ARTQHSLVLA RLRGRGRAHG RPCCQPTSVA DVTFLDDQHH WQQLPQLSAA ACGCGG
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using human TT medullary thyroid cancer cells is less than 0.1 ng/ml, corresponding to a specific activity of > 1.0 × 10 ⁷ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 30 % ACN, 0.1 % TFA, 150 mM NaCl.
Endotoxin:	Less than 0.1 EU/µg of rMuPersephin as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Murine Persephin

Persephin belongs to the GDNF ligand subfamily of the TGF-beta superfamily. It promotes the survival and growth of central dopaminergic and motor neurons, and also is involved in kidney development. But persephin does not support survival of peripheral neurons.