

**PrimeGene Technical Data Sheet**

---

<b>Catalog Number:</b>	224-24
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 10.3 kDa, a single, non-glycosylated polypeptide chain containing 93 amino acids.
<b>Quantity:</b>	5µg/20µg/1000µg
<b>AA Sequence:</b>	VTIPSSCCTS FISKKIPENR VVSYQLANGS ICPKAGVIFI TKKGHKICTD PKLLWVQRHI QKLDKKNQP SKGAKAVRTK FAVQRRRGNS TEV
<b>Purity:</b>	> 97 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using murine lymphocytes is in a concentration of 10-100 ng/ml.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
<b>Endotoxin:</b>	Less than 1 EU/µg of rMuEotaxin-2/CCL24 as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

***Murine Eotaxin-2/CCL24***

CCL24, also named MPIF-2, Eotaxin-2 and Ckβ6, is a novel CC chemokine recently identified. It is a secreted protein, encoded by CCL24 gene, and produced by activated monocytes and T lymphocytes. CCL24 signals through the CCR3 receptor and has functions of chemotactic activity for resting T-lymphocytes and eosinophils, but none for monocytes and activated lymphocytes. The plasma levels of CCL24 and the aspirin-exacerbated respiratory disease (such as asthma) morbidity rate have positive correlation.