

**Recombinant Rat Cytokine-induced Neutrophil
Chemoattractant 3/CXCL2
(rRtCINC-3/CXCL2)
PrimeGene Technical Data Sheet**

Catalog Number:	241-02
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 7.9 kDa, a single, non-glycosylated polypeptide chain containing 73 amino acids.
Quantity:	5µg/25µg/1000µg
AA Sequence:	VVVASELRCQ CLTTLPRVDF KNIQSLTVTP PGPHCAQTEV IATLKDGHVEV CLNPEAPLVQ RIVQKILNKG KAN
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using rat neutrophils is in a concentration range of 10-100 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rRtCINC-3/CXCL2 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Rat Cytokine-induced Neutrophil Chemoattractant 3/CXCL2

Rat CXCL2 also named CINC-3, belongs to the CXC chemokine family. It is encoded by the gene CXCL2 and shares 90 % amino acid sequence with CXCL1/GROα. It is member of the intercrine alpha (chemokine C-X-C) subfamily of chemokines. This chemokine is expressed by cytokine-stimulated rat alveolar macrophages and fibroblasts. The functional receptor for CXCL2 has been identified as CXCR2. CXCL2 is chemotactic for polymorphonuclear leukocytes and hematopoietic stem cells. Similar to other GRO proteins, CXCL2 is potent neutrophil attractants and activators. In addition, it is also active toward basophils. The amino acid sequence of rat CXCL2 is 88 % identical to murine CXCL2.