

**Recombinant Human High-Mobility
Group Box 1, His
(rHuHMGB1, His)
PrimeGene Technical Data Sheet**

Catalog Number:	601-06
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 26.0 kDa, a single non-glycosylated polypeptide chain containing 223 amino acids with 6 × His at C-terminus.
Quantity:	10µg/50µg/1000µg
AA Sequence:	MGKGDPPKPR GKMSYAFFV QTCREEHKKK HPDASVNFSE FSKKCSERWK TMSAKEKGGKF EDMAKADKAR YEREMKTYIP PKGETKKKFK DPNAPKRPPS AFFLFCSEYR PKIKGEHPGL SIGDVAKKLG EMWNNTAADD KQPYEKKA LKEKYEKDIA AYRAKGPDA AKKGVVKA EK SKKKKEEEED EDEEDEEEEE EDEEDEEEEE DDDDELEHHH HHH
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Data not available.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rHuHMGB1, His as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human High-Mobility Group Box 1

Human High-mobility group box 1 protein (HMGB1), previously known as HMG-1 or amphoterin, is a member of the high mobility group box family of non-histone chromosomal proteins. Human HMGB1 is expressed as a 30 kDa, 215 amino acid (a.a.) single chain polypeptide containing three domains: two N-terminal globular, 70 a.a. positively charged DNA-binding domains (HMG boxes A and B), and a negatively charged 30 a.a. C-terminal region that contains only Asp and Glu.4, 5 Residues 27 - 43 and 178 - 184 contain a NLS. Posttranslational modifications of the molecule have been reported, with acetylation occurring on as many as 17 lysine residues. HMGB1 is expressed at high levels in almost all cells. It was originally discovered as a nuclear protein that could bend DNA. Such bending stabilizes nucleosome formation and regulates the expression of select genes upon recruitment by DNA binding proteins.