

PrimeGene Technical Data Sheet

Catalog Number:	601-14
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 13.2 kDa, a homodimer consisting of two 60 amino acid polypeptide chains, which includes a 40-amino acid trefoil motif containing three conserved intramolecular disulfide bonds.
Quantity:	5µg/20µg/1000µg
AA Sequence:	EAQTETCTVA PRERQNCGFP GVTSPQCANK GCCFDDTVRG VPWCFYPNTI DVPPEEECEF
Purity:	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by a chemotaxis bioassay using human MCF-7 cells is less than 10 µg/ml, corresponding to a specific activity of > 100 IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Endotoxin:	Less than 1 EU/µg of rHuTFF1 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Trefoil Factor 1

Trefoil factor 1 belongs to the trefoil factor family that consists of three members named TFF1, TFF2 and TFF3. They are characterized by having at least one copy of the trefoil motif, a 40-amino acid domain that contains three conserved disulfides. The TFFs are stable secretory proteins expressed highly in the gastrointestinal tract (gastric mucosa). TFF1 is an essential protein for normal differentiation of the antral and pyloric gastric mucosa and functions as a stabilizer of the mucous gel overlying the gastrointestinal mucosa that provides a physical barrier against various noxious agents. It is studied because of commonly expressed in tumors as well.