

PrimeGene Technical Data Sheet

Catalog Number:	661-02
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 52.7 kDa, a single non-glycosylated polypeptide chain containing 503 amino acids, with Leu, Glu and 6 × His at C-terminus.
Quantity:	10μg/50μg/1000μg
AA Sequence:	MAQVINTNSL SLLTQNNLNK SQSALGTAIE RLSSGLRINS AKDDAAGQAI ANRFTANIKG LTQASRNAND GISIAQTTEG ALNEINNNLQ RVRELAVQSA NSTNSQSDLD SIQAEITQRL NEIDRVSGQT QFNGVKVLAQ DNTLTIQVGA NDGETIDIDL KQINSQTLGL DTLNVQQKYK VSDTAATVTG YADTTIALDN STFKASATGL GGTDQKIDGD LKFDDTTGKY YAKVTVTGGT GKDGYYEVSV DKTNGEVTLA GGATSPLTGG LPATATEDVK NVQVANADLT EAKAALTAAG VTGTASVVKM SYTDNNGKTI DGGLAVKVGD DYYSATQNKD GISISINTTKY TADDGTSKTA LNKLGGADGK TEVVSIGGKT YAASKAEGHN FKAQPDIAEA AATTENPLQ KIDAALAQVD TLRSDLGAVQ NRFNSAITNL GNTVNNL TSA RSRIEDSDYA TEVSNMSRAQ ILQQAGTSVL AQANQVPQNV LSLLRLEHHH HHH
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Data is not available.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/μg of rFlagellin, His as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Flagellin

Flagellin arranges itself in a hollow cylinder to form the filament in bacterial flagellum. It is the principal substituent of bacterial flagellum, and is present in large amounts on nearly all flagellated bacteria. Mammals often have acquired immune responses (T-cell and antibody responses) to flagellated bacterium and the plant defense mechanisms can be activated by the conserved N-terminal part of flagellin.